Department of Health

COVID-19 Update




Charlotte Update

35,466 total cases

- October 2021- 764 October 2020- 662
- November 2021- 339 November 2020- 1,387
- December 2021-1,346 December 2020-2,411
- January 2022- 7,770 January 2021- 2,086
- February 2022-2,334 February 2021-927
- March 2022-265 March 2021-24
- New weekly case Positivity 4.2%, State Average 2.4%
- Overall positivity for outbreak 24.3%; State 26.5%

Treatment Options



- Monoclonal site open at Tringali
 - M-Sat
 - 9a-5p
 - Accepting walk-ups
 - Will transition to appts once web page is updated.

https://floridahealthcovid19.gov/treatments/treatmentlocator/

Charlotte Status

Medium

What Prevention Steps Should You Take Based on Your COVID-19 Community Level?

Low	Medium	High
Stay up to date with COVID-19 vaccines Get tested if you have symptoms	 If you are at high risk for severe illness, talk to your healthcare provider about whether you need to wear a mask and take other precautions Stay up to date with COVID-19 vaccines Get tested if you have symptoms 	 Wear a mask indoors in public Stay up to date with COVID-19 vaccines Get tested if you have symptoms Additional precautions may be needed for people at high risk for severe illness

People may choose to mask at any time. People with symptoms, a positive test, or exposure to someone with COVID-19 should wear a mask. Masks are required on <u>public transportation</u> and may be required in other places by local or state authorities.

Strategy/Updates





COVID-19 vaccines are recommended for persons 5 years of age and older within the scope of the Emergency Use Authorization or Biologics License Application for the vaccine. The table below provides guidance for COVID-19 vaccination schedules based on age and medical condition. Detailed information can be found in CDC's Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Approved or Authorized in the United States (link below).

Table 1. Immunization schedule for persons 5 years of age and older

Recipient Product*†		Persons Who ARE NOT Moderately or Severely Immunocompromised		Persons Who ARE Moderately or Severely Immunocompromised	
		Primary Series ^{‡§}	Booster Dose ^{‡¶}	Primary Series ^{‡5}	Booster Dose ^{‡¶}
Type: mRNA	vaccine				
5–11 years	Pfizer-BioNTech Ages: 5–11 years Orange cap	2 doses. Separate: Dose 1 and 2 by at least 3 weeks **	Not recommended	3 doses. Separate: Dose 1 and 2 by at least 3 weeks. Dose 2 and 3 by at least 4 weeks.	Not recommended
12-17 years	Pfizer-BioNTech Ages: 12 years and older Gray cap or Purple cap	2 doses. Separate: Dose 1 and 2 by at least 3 - 8 weeks."	At least 5 months after Dose 2	3 doses. Separate: Dose 1 and 2 by at least 3 weeks. Dose 2 and 3 by at least 4 weeks.	At least 12 weeks after Dose 3
18 years and older	Pfizer-BioNTech Ages: 12 years and older Gray cap or Purple cap	2 doses. Separate: Dose 1 and 2 by at least 3 - 8 weeks.**	At least 5 months after Dose 2	3 doses. Separate: Dose 1 and 2 by at least 3 weeks. Dose 2 and 3 by at least 4 weeks.	At least 12 weeks after Dose 3
	Moderna	2 doses. Separate: Dose 1 and 2 by at least 4 - 8 weeks."*	At least 5 months after Dose 2	3 doses. Separate: Dose 1 and 2 by at least 4 weeks. Dose 2 and 3 by at least 4 weeks.	At least 12 weeks after Dose 3
Recipient Age	Product*†	Persons Who ARE NOT Moderately or Severely Immunocompromised		Persons Who ARE Moderately or Severely Immunocompromised	
		Primary Series†§	Booster Dose ^{‡¶}	Primary Series ^{‡§}	Booster Dose ^{‡¶}
Type: Viral v	vector vaccine				
18 years and older	Janssen ^{††}	1 dose	At least 8 weeks after Dose 1	2 doses. Separate: Dose 1 and 2 by at least 28 days ^{‡‡} Dose 2 MUST be a mRNA vaccine	At least 8 weeks after Dose 2

- CDC updates community level guidelines based upon transmission of COVID-19.
 - CDC County Check tool added to web.
- Three test sites in West, Central & South
- Multiple therapeutics available on state web page. (See Slide 7)
- Continue testing and vaccination strategies.
- Continue rapid Infection control assessments.
- Vaccination efforts continue. (75% total pop 5+)
- Continued focus on ILI and respiratory illnesses.
- Continue to monitor unmet needs and respond appropriately.
- Federal site for home tests. <u>https://www.covidtests.gov/</u>

CDC Guidance

When to Stay Home

Calculating Quarantine

The date of your exposure is considered day 0. Day 1 is the first full day after your last contact with a person who has had COVID-19. Stay home and away from other people for at least 5 days. Learn why CDC updated guidance for the

After quarantine

IF YOU Were exposed to COVID-19 and are NOT up-to-date on COVID-19 vaccinations

Quarantine for at least 5 days

Stay home Stay home and quarantine for at least 5

Wear a well-fitted mask if you must be around others in your home.

Do not travel.

No quarantine

No quarantine

symptoms.

Get tested

Even if you don't develop symptoms, get tested at least 5 days after you last had close contact with someone with COVID-19.

You do not need to stay

home unless you develop

Even if you don't develop

symptoms, get tested at

had close contact with

least 5 days after you last

someone with COVID-19.

You do not need to stay

home unless you develop

IE VOLL Were exposed to COVID-19 and are up-todate on COVID-19 vaccinations

IF YOU were exposed to COVID-19 had confirmed COVID-19 within the past 90 days (you tested positive using a viral test)

Watch for symptoms until 10 days after you last had close contact with someone with COVID-19.

Avoid travel

It is best to avoid travel until a full 10 days after you last had close contact with someone with COVID-19.

If you develop symptoms Isolate immediately and get tested. Continue to stay home until you know the results. Wear a wellfitted mask around others

Watch for symptoms Watch for symptoms until 10 days after you last had close contact with someone with COVID-19.

If you develop symptoms Isolate immediately and get tested. Continue to stay home until you know the results. Wear a wellfitted mask around others

Watch for symptoms Watch for symptoms until 10 days after you last had close contact with someone with COVID-19

If you develop symptoms Isolate immediately and get tested. Continue to stay home until you know the results. Wear a wellfitted mask around

Take precautions until day 10 Wear a mask

Watch for symptoms

Wear a well-fitted mask for 10 full days any time you are around others inside your home or in public. Do not go to places where you are unable to wear a mask.

If you must travel during days 6-10. take precautions.

Avoid being around people who are at high risk

Take precautions until day 10

Wear a mask

Wear a well-fitted mask for 10 full days any time you are around others inside your home or in public. Do not go to places where you are unable to wear a mask.

Take precautions if traveling

Avoid being around people who are at high risk

Take precautions until day 10

Wear a well-fitted mask for 10 full days any time you are around others inside your home or in public. Do not go to places where you are unable to wear a mask.

Take precautions if traveling

Avoid being around people who are at high risk

Calculating Isolation

Day 0 is your first day of symptoms or a positive viral test. Day 1 is the first full day after your symptoms developed or your test specimen was collected. If you have COVID-19 or have symptoms, isolate for at least 5 days.

IF YOU Tested positive for COVID-19 or have symptoms, regardless of vaccination status

Stay home for at least 5 davs Stay home for 5 days and isolate from others in your home.

Wear a well-fitted mask if you must be around others in your home.

Do not travel.

Ending isolation if you had symptoms End isolation after 5 full days if you are fever-free for 24 hours (without the use of fever-reducing medication) and your

symptoms are improving.

Ending isolation if you did NOT have symptoms End isolation after at least 5 full days after your positive test.

If you were severely ill with COVID-19 or are immunocompromised You should isolate for at least 10 days. Consult your doctor before ending isolation.

Take precautions until day 10

Wear a mask

Wear a well-fitted mask for 10 full days any time you are around others inside your home or in public. Do not go to places where you are unable to wear a mask.

Do not travel

Do not travel until a full 10 days after your symptoms started or the date your positive test was taken if you had no symptoms.

Avoid being around people who are at high risk

CDC Healthcare Worker Guidance

Work Restrictions for HCP With SARS-CoV-2 Infection and Exposures

"Up to Date" with all recommended COVID-19 vaccine doses is defined in Stay Up to Date with Your Vaccines I CDC

For more details, including recommendations for healthcare personnel who are iimmunocompromised, have severe to critical illness, or are within 90 days of prior infection, refer to Interim Guidance for Managing Healthcare Personnel with SARS-CoV-2 Infection or Exposure to SARS-CoV-2 (conventional standards) and Strategies to Mitigate Healthcare Personnel Staffing Shortages (contingency and crisis standards).

Work Restrictions for HCP With SARS-CoV-2 Infection

Vaccination Status	Conventional	Contingency	Crisis
Up to Date and Not Up to Date	10 days OR 7 days with negative test ¹ , if asymptomatic or mild to moderate illness (with improving symptoms)	5 days with/without negative test, if asymptomatic or mild to moderate illness (with improving symptoms)	No work restriction, with prioritization considerations (e.g., types of patients they care for)

Work Restrictions for Asymptomatic HCP with SARS-CoV-2 Exposures

Vaccination Status	Conventional	Contingency	Crisis
Up to Date	No work restrictions, with negative test on days 1 ⁺ and 5–7	No work restriction	No work restriction
Not Up to Date	10 days OR 7 days with negative test*	No work restriction with negative tests on days 1*, 2, 3, & 5–7 (if shortage of tests prioritize Day 1 to 2 and 5-7)	No work restrictions (test if possible)

†Negative test result within 48 hours before returning to work ‡For calculating day of test: 1) for those with infection consider day of symptom onset (or first positive test if asymptomatic) as day 0; 2) for those with exposure consider day of exposure as day 0



CHRONAL CONTROL

cdc.gov/coronavirus